


SCIP - INPATIENT VTE PROPHYLAXIS OPTIONS FOR SURGERY



Applicable to discharges 1Q14 (01-01-2014) through 3Q14 (09-30-2014)

2008 American College of Chest Physicians

INTRACRANIAL NEUROSURGERY

Any of the following:

- Intermittent pneumatic compression devices (IPC) with or without graduated compression stockings (GCS)
- Low-dose unfractionated heparin (LDUH)
- Low molecular weight heparin (LMWH)
- LDUH or LMWH combined with IPC or GCS

GENERAL SURGERY

Any of the following:

- Low-dose unfractionated heparin (LDUH)
- Low molecular weight heparin (LMWH)
- Factor Xa Inhibitor
- Intermittent pneumatic compression devices (IPC)

GYNECOLOGIC SURGERY

Any of the following:

- Low-dose unfractionated heparin (LDUH)
- Low molecular weight heparin (LMWH)
- Factor Xa Inhibitor
- Intermittent pneumatic compression devices (IPC)
- LDUH or LMWH or Factor Xa Inhibitor combined with IPC or GCS

UROLOGIC SURGERY

Any of the following:

- Low-dose unfractionated heparin (LDUH)
- Low molecular weight heparin (LMWH)
- Factor Xa Inhibitor
- Intermittent pneumatic compression devices (IPC)
- LDUH or LMWH or Factor Xa Inhibitor combined with IPC or GCS

ELECTIVE TOTAL KNEE OR TOTAL HIP REPLACEMENT

Any of the following:

- Low molecular weight heparin (LMWH)
- Factor Xa Inhibitor
- Oral Factor Xa Inhibitor*
- Warfarin
- Intermittent pneumatic compression devices (IPC)
- Venous foot pump (VFP)
- Low-dose unfractionated heparin (LDUH)
- Aspirin

HIP FRACTURE SURGERY

Any of the following:

- Low-dose unfractionated heparin (LDUH)
- Low molecular weight heparin (LMWH)
- Factor Xa Inhibitor
- Warfarin
- Intermittent pneumatic compression devices (IPC)
- Aspirin

*The U.S. Food and Drug Administration has approved Xarelto (rivaroxaban) to reduce the risk of blood clots, deep vein thrombosis (DVT) and pulmonary embolism (PE) following knee or hip replacement surgery **ONLY**.